

Biochemical Studies On The Effect Of Ponceau 4R And/Or Vitamin E Treatment On Young Male Albino Rats

*Eman Helal; *Al-Kassas, M; **Ghada Z A Soliman; *Zahkok, S;
*Abdel Wahed, H.

*Zoology Dep. Faculty of Science, Al-Azhar University (Girls),
**National Nutrition Institute, Cairo, Egypt;

Abstract

Introduction: Due to now a day's high technology, which make foods more attractable to children. This study aims to study the effect of Ponceau 4R (food colorant) and/or vitamin E (antioxidant) on rats.

Material And Methods: Certain parameters were measured as percentage of body weight change, body temperature, heart beat, Red & white blood cells count, hemoglobin concentration, hematocrite value, serum total lipids, serum cholesterol, serum total protein, serum albumin, globulin, A/G ratio, serum glucose, serum alanine transaminase and aspartate transaminase (ALT & AST) activity and serum total cholinesterase. The organs (brain, liver, kidney, heart and skeletal muscle) were removed, cleared then weighted. Organs were prepared for total protein, total lipid determination and other biochemical analysis.

Results: Body weight, respiration rate was significantly decreased till the 3rd month followed by a significant increase till the 6th month. WBCs count; serum total lipids, liver total lipids, serum glucose, serum cholinesterase activity were significantly reduced. Muscle total lipids, serum total cholesterol, serum total protein, serum total albumin, serum AST activity, liver AST activity were significantly increased.

Conclusion: Due to the hazardous effect of food colourants as Ponceau 4R, it is recommended that the use of Ponceau 4R as food colourant must be limited and use vitamin E as antioxidant to prevent the toxic effect of food colouring substance.

Key Words: Ponceau 4R, Rats, Vitamin E

Introduction

Food additives are substances that are not normally consumed as food by itself or considered as food ingredient. Whether it has nutritive value or not, its addition to foods during manufacture must be according to FAO & WHO (1991) regulations. Antioxidant is a molecule stable enough to donate an electron to trap the free radical and neutralize it. Vitamin E is natural antioxidant present in foods and stored mainly in adipose tissue (Bagchi and Puri, 1998). It protects polyunsaturated fatty acids in cell membrane from peroxidation, it metabolize free radicals and reduces the risk of disease outcomes and plays a role in cancer prevention and control (Bagchi and Puri 1998; and Metin *et al.*, 2002). Ponceau 4R is food colourant that is used all over the world in great amounts. This colourant (Ponceau 4R) was

evaluated for acceptable daily intake by the Joint FAO/WHO Expert Committee on Food Additives (Bär & Grienpentrog 1960; Gaunt *et al.*, 1967; and Stevenson *et al.*, 1980, 1981). Since the previous evaluation, additional data have become available. Metabolism studies indicate that Ponceau 4R is rapidly absorbed from the gastrointestinal tract and undergoes extensive metabolism with the formation of naphthionic acid and 7-hydroxy-8-amino-naphthalene-1,3-disulfonic acid. No marked accumulation of radioactivity in any tissue was found following a single dose of ¹⁴C-labelled colouring to untreated rats, mice and guinea pigs or rats given repeated doses of unlabelled colouring for 28 days. Ponceau 4R consists essentially of trisodium-1- (4-sulpho-1- naphthylazo) 2-naphthol-6, 8 disulfonate. It is also known

as Food Red 7, C. I. 16255, Cochineal Red A, New Coccine, Acid Red 18, SX purple, and E 124. Ponceau 4R is a synthetic red azo dye which can be used in a variety of food products. Since it is an azo dye, it may elicit intolerance in people allergic to salicylates (aspirin). Additionally, it is a histamine liberator, and may intensify symptoms of asthma. Ponceau 4R is considered carcinogenic in some countries, including the USA, Norway, and Finland, and it is currently listed as a banned substance by the U.S. Food and Drug Administration (FDA)¹, although it is still used in developing countries.

Aim Of The Work

Due to the increasing rate of consuming foods containing food colorants (as Ponceau 4R) especially by children with no control on them, this work aims to study its effect on rats lipid profile, kidney and liver function, and some enzymes and to see if presence of the natural antioxidant as vitamin E can minimize the hazards effect and protect our children.

Material And Methods

The present study was carried out on forty immature growing male albino rats of about one month age. The weight range was from 40-50 g. The animals were kept in clean cages. Food and water were added *ad libitum*. The Experimental animals were divided into four groups (10 rats/group) as follows: Group 1 (G 1): serve as control group; Group 2 (G 2): supplemented with vitamin E (150 mg/kg BW); Group 3 (G 3): supplemented with Ponceau 4R (0.19 mg/kg BW); Group 4 (G 4): supplemented with Ponceau 4R and vitamin E. These treatments were administered orally by stomach tube for six months. All animal groups were weighted before beginning and monthly during the experiment period. Half of each group was decapitated after 3 month, while the other half were left till the end of 6 months. At the end of the experiment, animals were decapitated and

blood samples were taken in dry clean centrifuge tube. Serum was separated and kept at -20 °C until analysis. Blood samples for hematological analysis were taken by heparinized capillary tube. The organs (brain, liver, kidney, heart and skeletal muscle) were removed and cleared in isotonic saline solution then weighted. A piece of each organ were weighted and put in appropriate amount of 30 % KOH for total protein determination and in conc. H₂SO₄ for total lipid determination. Other pieces of organ were weight and put in saline solution then homogenized for biochemical analysis.

Measured Parameters

Certain parameters were measured. Percentage of body weight change was recorded monthly. Heart rates and respiration rate were recorded according to the method of Soliman *et al.* (1973). Red & white blood cells were counted according to Mitruka *et al.* (1977). Hemoglobin level was determined according to Van Kampen and Zijlstra (1961). Hematocrite value was determined according to Rodak (1995). Serum total lipids were determined according to Knight *et al.* (1972). Serum cholesterol was determined according to Martinek *et al.* (1970). Serum total protein was determined according to Doumas (1975). Serum albumin was determined according to Doumas *et al.* (1971). Serum glucose was determined according to Trinder (1969). Serum alanine transaminase and aspartate transaminase (ALT & AST) activity were determined according to Reitman & Frankel (1957). Serum total cholinesterase was determined according to Gorun *et al.* (1978).

Statistical Analysis

Data are expressed as Mean \pm SE. Data were assessed by t-test (Armitage 1974; and Lenter *et al.*, 1982). P-values < 0.05 were considered statistically significant.

Results

Table (1) shows percentage of body weigh, heart rate, rectal temperature and respiration rate changes of male albino rats treated with Ponceau 4R and vitamin E for

¹ "Artificial Food Colouring Warning", BBC News, May 8, 2007.

three and six months. Data of table (1) shows insignificant change in body weight, heart rate, rectal temperature in rats treated with Ponceau 4R and/or vitamin E in comparison with the control group. Also data of table (1) revealed that a significant decrease ($P < 0.05$) in respiration rate was recorded in rats treated with Ponceau 4R after 3 months and the opposite was noticed after 6 months where a significant increase ($P < 0.05$) was recorded when compared with the control group. Rats treated with vitamin E alone or in combination with Ponceau 4R showed insignificant differences in respiration rate when compared with the control group during the experimental period.

Data of table (2) shows insignificant change in the percentage of organ/body weight changes of male albino rats treated with Ponceau 4R and/or vitamin E for three and six months in brain, kidney and heart. On the other hand, a highly significant decrease ($P < 0.01$) was recorded in hepatosomatic index in rats treated with Ponceau 4R after 3 & 6 months when compared with the control group.

Data of table (3) shows insignificant change in the red blood corpuscles (RBCs) count, hemoglobin (Hb) concentration and hematocrite (Hct) value of rats treated with Ponceau 4R and/or vitamin E for three and six months. Also table (3) revealed non significant difference in white blood corpuscles (WBCs) count in all treated groups after 3 & 6 months in comparison with the control group except rats treated with Ponceau 4R where they showed a significant decrease ($P < 0.05$) after 6 months.

From table (4), it was clear that treating rats with vitamin E, combination of vitamin E and Ponceau 4R revealed insignificant difference in total lipids, total cholesterol, total protein, albumin, globulin levels after 3 & 6 months of treatment. On the other hand, a significant decrease ($P < 0.01$) in total lipid, total protein, albumin and globulin level was observed in rats treated with Ponceau 4R (only) after 3 & 6 months of treatment in comparison with the control group, while rats treated with Ponceau 4 R alone after 3 & 6 months of treatment showed a significant increase (P

< 0.01) in total cholesterol level in comparison with the control group.

Table (5) shows tissue total lipids (mg/g tissue) levels of male albino rats in organs as brain, liver, kidney, muscle and heart treated with Ponceau 4R and vitamin E for three and six months. It can be seen that rats supplemented with vitamin E showed insignificant difference in total lipid content in brain, liver, muscle, kidney and heart tissues after 3 & 6 months of treatment in comparison with the control group. Rats treated with Ponceau 4R showed a significant decrease ($P < 0.05$) in total lipids of liver tissue, while a significant increase ($P < 0.01$) in total lipids of muscle tissue was observed. Also rats treated with Ponceau 4R combined with vitamin E revealed a significant increase ($P < 0.05$) in total lipids in muscle tissue in comparison with the control group after 3 & 6 months.

Data of table (6) shows that rats treated with Ponceau 4R and/or vitamin E revealed insignificant change in total cholesterol in brain, liver, kidney, muscle and heart tissues after three and six months in comparison with the control group.

Table (7) shows tissue total protein (mg/g tissue) of different organs of male albino rats treated with Ponceau 4R and/or vitamin E for three and six months. Data of table (7) shows that rats treated with Ponceau 4R and/or vitamin E revealed insignificant change in total protein in brain, liver, kidney, muscle and heart tissues after three and six months in comparison with the control group.

Table (8) shows serum glucose, AST, ALT and cholinesterase of male albino rats treated with Ponceau 4R and vitamin E for three and six months. Data of table (8) revealed that rats treated with vitamin E or with dual treatment Ponceau 4R and vitamin E showed insignificant change in serum glucose, AST, ALT and cholinesterase level after 3 & 6 months. Rats treated with Ponceau 4R alone showed significant decrease in glucose level after 3 & 6 months ($P < 0.05$). Serum activity of AST of rats treated with Ponceau 4R showed a significant increase ($P < 0.05$) after 3 & 6 months when compared with the control group. Rats treated with Ponceau

4R alone showed insignificant change in ALT activity after 3 & 6 months. Obtained data (in table 8) revealed insignificant difference in cholinesterase activity in all treated groups after 3 & 6 months in comparison with the control group, except rats treated with Ponceau 4R, which showed highly significant, decrease ($P < 0.01$) in cholinesterase activity after 3 & 6 months.

Table (9) shows tissue AST & ALT activity (U/g tissue) of different organs of male albino rats treated with Ponceau 4R and vitamin E for three and six months. It is clear from the table that rats treated with

vitamin E alone or in combination with Ponceau 4R or treated with Ponceau 4R alone revealed insignificant change in AST, ALT activity in brain, liver, kidney, muscle, and heart tissues after 3 & 6 months when compared with the control group except AST activity of rats liver treated with Ponceau 4R alone where it showed a significant increase ($P < 0.05$). Data of the table (9) reveal that despite the insignificant change of AST activity in brain, but the activity is increased insignificantly especially with rats treated with vitamin E alone (41.37 %).

Table (1): Percentage of body weight, heart beat rate, rectal temperature and respiration rate changes of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	3 Months				6 Months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Body Weight	23.56±2.83	25.13±1.30	27.36±2.14	25.39±0.93	22.11±1.27	19.48±1.72	16.42±2.84	20.91±1.59
Heart Beat Rate	315.0±4.01	311.0±3.48	318.0±3.27	314.0±3.73	312.0±3.74	316.0±4.00	310.0±3.16	314.0±4.31
Rectal Temperature	37.35±0.04	37.41±0.07	37.40±0.06	37.42±0.05	37.26±0.05	37.24±0.08	37.27±0.06	37.37±0.08
Respiration Rate	110.6±1.82	115.3±1.59	103.1±1.93*	113.6±1.49	109.4±2.79	113.2±1.96	120.4±3.2*	114.2±2.33

*: Significant, $P < 0.05$

Table (2): Percentage of organ /body weight changes of male albino rats treated with Ponceau 4R and vitamin E for three and six months

	3 months				6 months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Brain	0.95±0.06	1.03±0.12	0.75±0.08	0.78±0.08	0.59±0.01	0.63±0.04	0.58±0.03	0.63±0.03
Liver	3.43±0.10	3.36±0.09	2.90±0.04**	2.99±0.18	3.01±0.08	2.85±0.09	2.61±0.051**	2.92±0.07
Kidney	0.62±0.02	0.64±0.04	0.53±0.04	0.57±0.01	0.65±0.04	0.63±0.04	0.62±0.02	0.56±0.02
Heart	0.34±0.03	0.51±0.18	0.31±0.01	0.29±0.01	0.29±0.01	0.31±0.02	0.33±0.02	0.34±0.03

** : Highly Significant, $P < 0.01$

Table (3): Mean values of red blood corpuscles (RBCs) count, white blood corpuscles (WBCs) count, hemoglobin (Hb) concentration and hematocrite (Hct) value male albino rats treated with Ponceau 4R and vitamin E for three and four months.

	3 months				6 months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
RBCs	7.88±0.12	8.06±0.25	7.40±0.18	7.90±0.29	7.50±0.41	6.82±0.31	6.52±0.14	6.74±0.37
WBCs	11.56±0.88	9.90±0.34	8.12±0.78	10.20±0.34	12.32±0.40	11.38±0.24	10.34±0.49*	12.14±0.25
Hb g/dl	15.18±0.59	15.80±0.25	13.25±0.80	16.18±0.22	14.38±0.38	14.40±0.27	12.34±0.86	14.72±0.64
Hct %	47.06±1.93	50.24±1.10	47.72±0.93	50.56±1.67	45.76±1.37	44.62±2.39	42.58±2.17	49.42±1.36

** : Highly Significant, $P < 0.01$

Table (4): Serum total lipids, total cholesterol, total protein, albumin, globulin and A/G ratio of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	3 Months				6 Months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Total Lipids (mg/dl)	352.32±8.21	352.54±12.3	262.7±5.12**	334.36±14.53	357.52±8.05	358.94±8.52	311.78±5.04**	338.78±7.13
Total Cholesterol (mg/dl)	123.24±2.01	122.32±1.54	145.8±5.37**	127.06±3.32	114.34±4.91	118.76±5.81	133.0±2.45**	115.92±2.91
Total Protein (g/dl)	6.90±0.07	6.98±0.12	5.62±0.12**	6.76±0.24	6.76±0.10	6.56±0.13	5.76±0.10**	6.56±0.14
Albumin (g/dl)	4.50±0.08	4.16±0.19	3.60±0.07**	4.25±0.15	4.43±0.07	4.64±0.13	3.64±0.09**	4.48±0.07
Globulin (g/dl)	2.40±0.14	2.28±0.10	2.02±0.07*	2.51±0.14	2.33±0.04	1.92±0.21	2.12±0.16	2.08±0.17
A/G ratio	1.88±0.14	1.48±0.18	1.78±0.05	1.69±0.16	1.90±0.02	2.42±0.34	1.72±0.17	2.15±0.21

*: Significant, P < 0.05, **: Highly Significant, P < 0.01

Table (5): Tissue total lipids (mg/g tissue) of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	3 Months				6 Months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Brain	79.48±5.63	89.58±3.88	74.86±3.20	76.12±2.89	59.44±4.28	69.84±2.97	62.72±2.27	53.94±3.19
Liver	91.76±3.51	96.10±5.93	74.74±6.41*	77.62±7.46	86.08±4.29	75.50±3.69	65.68±5.47*	73.62±4.44
Kidney	55.92±1.90	60.08±1.00	52.34±1.74	58.58±4.16	62.82±3.61	56.90±2.30	59.84±2.84	59.52±3.10
Muscle	51.08±6.74	65.60±4.30	80.10±2.41**	73.90±3.07**	40.98±2.30	47.24±4.62	58.40±1.98**	51.40±3.00**
Heart	58.60±2.61	63.62±2.88	62.90±5.69	64.36±2.08	53.58±6.50	51.66±3.30	55.90±2.71	45.56±3.83

*: Significant, P < 0.05, **: Highly Significant, P < 0.01

Table (6): Tissue total cholesterol (mg/g tissue) of different organs of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	3 Months				6 Months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Brain	66.42±3.24	79.58±5.64	67.17±6.20	60.95±4.27	80.54±3.89	87.67±3.73	74.62±1.90	72.55±3.72
Liver	55.47±2.77	58.84±3.46	59.84±2.49	58.48±3.32	59.39±3.69	60.94±4.73	64.65±5.23	61.25±2.85
Kidney	61.33±2.72	62.43±3.83	62.79±3.91	58.11±3.65	61.56±2.46	63.54±2.22	66.08±2.76	59.94±2.04
Muscle	55.50±3.79	59.60±2.82	62.56±2.61	49.92±2.93	55.97±4.83	57.16±2.06	59.74±3.31	53.11±2.18
Heart	64.20±3.11	63.79±2.66	63.16±2.48	58.17±2.27	68.08±2.49	66.58±2.09	66.26±2.39	65.10±2.31

Table (7): Tissue total protein (mg/g tissue) of different organs of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	3 Months				6 Months			
	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E	Control	Vitamin E (150 mg/kg)	Ponceau 4R (0.19 mg/kg)	Ponceau 4R & Vitamin E
Brain	43.36±1.80	41.28±2.02	38.54±1.50	39.14±1.86	44.06±2.75	39.70±1.80	39.40±3.47	40.44±3.06
Liver	40.06±3.26	33.76±2.89	35.56±2.97	50.28±3.89	49.74±2.81	41.96±2.71	41.88±2.16	44.94±2.79
Kidney	26.04±3.77	29.20±3.32	22.18±2.69	22.20±1.99	44.22±4.44	42.78±3.94	39.34±2.15	42.18±2.38
Muscle	32.62±3.14	36.04±4.36	29.76±1.96	42.16±3.16	42.76±4.30	42.52±3.59	47.32±2.65	39.86±3.83
Heart	33.40±4.44	27.44±1.83	25.84±2.05	27.74±2.59	38.60±2.21	42.62±2.50	43.10±2.43	46.10±2.49

Table (8): Serum glucose, AST, ALT and cholinesterase of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

	Control	3 Months			Control	6 Months		
		Vitamin E 150 mg/kg	Ponccan 4R (0.19 mg/kg)	Ponccau 4R + vitamin E		Vitamin E 150 mg/kg	Ponccan 4R 0.19 mg/kg	Ponccau 4R + vitamin E
Glucose (mg/dl)	99.38±6.34	100.10±8.46	76.4±3.27**	114.30±3.20	96.34±4.16	90.60±3.64	79.60±3.81*	97.22±4.47
AST (U/l)	27.94±1.55	29.04±0.96	40.6±4.12**	30.20±2.00	30.98±0.40	28.38±1.23	35.40±1.47*	32.64±1.21
ALT (U/L)	24.50±1.50	26.30±1.01	23.30±2.78	25.98±1.14	24.64±1.19	25.50±1.32	28.34±1.24	25.23±0.84
Cholinesterase	11.70±0.89	9.22±1.19	7.96±0.44**	12.18±0.44	10.50±1.05	9.62±0.74	6.12±0.58**	10.46±0.36

*: Significant, $P < 0.05$, **: Highly Significant, $P < 0.01$

Table (9): Tissue AST & ALT (u/g tissue) of different organs of male albino rats treated with Ponceau 4R and vitamin E for three and six months.

		3 Months				6 Months			
		Control	Vitamin E 150 mg/kg	Ponccan 4R 0.19 mg/kg	Ponccau 4R + vitamin E	Control	Vitamin E 150 mg/kg	Ponccan 4R 0.19 mg/kg	Ponccau 4R + vitamin E
AST	Brain	17.84±1.92	25.22±2.32	21.40±2.90	22.30±1.91	22.30±1.88	23.02±2.14	24.66±1.10	23.12±2.21
	Liver	20.22±2.49	23.82±2.91	29.82±2.46*	20.60±2.24	21.72±2.61	22.02±1.86	30.62±2.31	19.78±1.67
	Kidney	20.60±2.27	22.90±3.52	23.12±2.13	19.24±2.41	21.12±1.38	18.24±1.74	19.56±1.80	18.68±1.61
	Muscle	18.62±1.64	20.26±2.10	20.70±2.23	20.82±2.64	18.54±1.68	17.76±1.26	17.32±1.35	18.16±2.25
	Heart	16.52±1.82	16.10±1.33	19.30±3.22	18.24±2.49	22.88±2.23	19.40±2.42	22.30±1.88	19.00±2.50
ALT	Brain	16.14±1.36	19.18±1.83	16.04±1.47	16.80±2.00	15.94±0.90	15.48±1.08	15.32±0.95	13.88±0.38
	Liver	14.40±1.48	20.78±2.03	15.28±1.35	13.56±1.60	15.44±1.12	16.24±1.32	16.08±1.23	16.18±1.36
	Kidney	15.22±1.77	21.54±3.44	15.32±1.80	18.92±2.97	16.66±1.13	16.52±1.17	15.78±1.40	16.84±1.37
	Muscle	13.70±1.02	15.34±1.80	15.48±1.36	15.94±2.25	19.98±2.55	16.18±1.33	18.08±1.16	16.24±1.32
	Heart	16.20±1.33	22.36±2.15	21.94±2.23	18.42±2.24	17.84±1.52	18.92±2.69	14.60±0.45	15.52±0.86

*: Significant, $P < 0.05$

Discussion

The present study is concerned with the effects of synthetic food colorant (Ponceau 4R) as preservative added to foodstuffs. Most of those that are used in developing countries are not permissible (AL-Sharkawi *et al.*, 1996) although now the regulation is towards using antioxidant.

Data of the present study showed that rats treated with vitamin E showed no significant change in body weight, which is in agreement with Meyer *et al.* (1989); and Eder and Kirchgessner (1998) who observed that the level of vitamin E in the diet had no effect on body weight gain. Data of the present study showed that rats treated with Ponceau 4R showed no significant change in body weight, which is in agreement with Brantom *et al.* (1987 a, b) who found no change in body weight of rats fed Ponceau 4R at dietary level (0, 50, 500, 1250 mg/kg BW), but they also found that high dose (toxic level) of Ponceau 4R lower body weight gain without any reduction in food intake. The reduction in body weight may be due to a reduction in food utilization (Grant and Butler 1989) or

vitamin C deficiency according to Uchida *et al.* (1990).

The present study revealed insignificant change in rats' heart rate in any treated groups till the end of the experiment. The stability of body temperature may be due to the high ability of rats to adjust their body temperature (Helal *et al.*, 1997). Data of the present study showed insignificant change in rats respiration rate in groups treated with vitamin E alone or with Ponceau 4R. On the other hand, groups treated with Ponceau 4R showed a significant decrease in respiration till the third month then a significant increase was noticed from the fourth month till the end of the experiment. This increment may be due to more oxygen requirement in the internal organs and it also attributed to the decrease in the red blood cell count and Hb content (Helal *et al.*, 1997). These are in disagreement with Mcdoniel *et al.* (1993) and Isliker *et al.* (1997) who suggested that supplementation with vitamin E stabilizes the cardiomyocyte as well as the mitochondria

membranes probably by preventing the detachment of their outer membrane. The inner mitochondria membrane is not sensitive to vitamin E depletion since the respiration rate was not affected Isliker *et al.* (1997).

Data in the present study indicate insignificant difference in percentage of heart, brain and kidney organs weight/body weight in all treated groups (vitamin E and/or Ponceau 4R), while hepatosomatic index showed a significant decrease in rats treated with Ponceau 4R. Farag *et al.* (1991) reported no change in the weight of liver, heart, kidney, spleen and brain organs in animals receiving natural antioxidant. They attributed this to the reason that stimulation has not taken place for enzymes necessary to breakdown these antioxidants.

The present work showed insignificant change in red blood cell count, Hb concentration and hematocrite value in groups treated with vitamin E. Moriguchi (1990) found that red blood corpuscles count increased with increasing content of vitamin E. The feeding of Ponceau 4R led to a significant decrease in erythrocyte count and hemoglobin level due to iron deficiency. Also the decrease may be due to the prevention of RBC synthesis via inhibition of erythropoiesis in bone marrow or may be due to iron deficiency anemia, hemolytic anemia, microcytic anemia. These results are in agreement with Mason *et al.* (1974); and Abu El-Zahab *et al.* (1997). The feeding of Ponceau 4R (200 mg/kg BW & 200 mg/L drinking water) led to a significant decrease in erythrocyte counts by 14.2 % (Gautam *et al.* 1986). The feeding of Ponceau 4R led to a significant decrease in WBC counts till the end of the experiment, while rats treated with vitamin E showed insignificant difference. Our results are in agreement with Tan *et al.* (1992) and Soheir *et al.* (1996). The reduction in WBCs count lower the defense mechanism (immune system) which play an important role in attacking and interacting with foreign antigens and initiating a primary immune response.

Data in the present study indicate insignificant change in serum total lipids of rats treated with vitamin E alone or in combination with Ponceau 4R, while treatment with Ponceau 4R alone revealed a

highly significant decrease in total lipids in serum, liver. These decreases in total lipids may be due to lipolysis, via stimulation of hormone sensitive lipase (Abd El-Dayem, 2002).

Data in the present study revealed a highly significant increase in serum total cholesterol levels of rats treated with Ponceau 4R till the end of the experiment. A highly significant decrease was observed in heart total cholesterol. The elevation in serum total cholesterol level may be attributed to the blockage of liver bile ducts, causing reduction or cessation of its secretion to the duodenum. Consequently, it appeared in the serum causing cholestasis (Hassan *et al.*, 1995 and Helal *et al.*, 1997). Also the elevation may be due to the mobilization of the free fatty acids from the adipose tissue to the blood stream and increase level of acetyl Co-A, leading to increase in the synthesis of cholesterol. The results are in agreement with the finding of Gomaa (1995); Hassan *et al.* (1995) and Helal *et al.* (1997).

The present study indicate a highly significant decrease in serum total proteins and albumin levels through out the experimental period in rats treated with Ponceau 4R. This reduction may be due to substantially of protein synthesis by the liver, this depression may be due to an alteration in the intracellular protein synthesis mechanism and that the oxidative enzyme changes were probably secondary in altering proteins. These results are in agreement with Hurkat (1977); Shakoory *et al.* (1988). Also Mekawy *et al.* (1988) and Amr *et al.* (1994) who reported that the decreases of protein might be due to reduction of serum globulin level supports with the disturbances on the immunoglobulin production, these was accompanied by a decrease of body weight gain and this may be a result of toxicity especially on the muscle. The decrease in serum albumin level may be due to loss of protein from the elementary tract, or due to decrease protein formation in the liver (impaired ability of the liver to form albumin). This result were in harmony with that obtained by Said *et al.* (1992), who concluded that the decrease in serum albumin level was due to trap of protein from the elementary tract or due to hepatic

necrosis. Rats treated with Ponceau 4R showed significant decrease in globulin level after 3 months of treatment, which may be due to the disturbance on immunoglobulin production.

Glucose is the key molecule in carbohydrate metabolism. The present data showed significant decrease in serum glucose level of rats treated with Ponceau 4R. This may be due to the effect of Ponceau 4R on cells of pancreas which lead to increase secretion of insulin hormone which reduce glucose in serum or the decrease may be due to liver disease, or due to adrenocortical insufficiency, anterior pituitary insufficiency and hypothyroidism as indicated by Katzung (1995).

The present data showed highly significant increase in serum AST activity of rats treated with Ponceau 4R through out the experiment. Elevation of transferase activity in blood has been considered as an indicator of tissue damage. However, other factors are considered for this process such as alteration in permeability of cell membrane, increasing the synthesis of the enzyme or decreasing the rate of degradation of the enzyme (Dinman *et al.* 1963; Luckens and Phelps 1969). Ignatov (1976) recorded that the elevation in serum AST is due to the degradation and necrosis of liver cells, which is accompanied by a damage of cell wall, cytolysis and so pouring a considerable amount of these mitochondrial enzymes into the blood stream. The present results are in agreement with those obtained by Yani *et al.* (1991); Abd El-Dayem (1992); and Ahmed & Mona (2000).

Generally the activities of ALT & AST are considerably increased following the administration of various hepatotoxic compounds, which lead to acute hepatocellular damage or extrahepatic obstructions by being considered as highly sensitive liver markers (Abu El-Zahab *et al.* 1997). Saleh (1986) related the elevation of ALT & AST activities to the damage of the liver cell, while Abu El-Zahab *et al.* (1997) report that coloring agents induce liver tissue damage and cause a significant increase of ALT, AST & alkaline phosphatase activities in rats serum.

Acetyl cholinesterase inhibition is one of the most important negative effects and

the interaction with the enzyme results in acute cholinergic poisoning (Wills 1972; and Howard & Janice 1989). Data of the present work showed that rats treated with Ponceau 4R showed a significant decrease in serum cholinesterase activity after 3 & 6 months of treatment. These decreases may be a sign of metabolic alterations of brain cholinergic synapses, which form a part of an inhibitory mechanism controlling the activity of cholinergic neurons. The inhibitory effect of Ponceau 4R administration on cholinesterase activity could be due to a direct action on the enzyme activity (Abd El-Rahiem *et al.* 1999) or due to hepatic parenchyma disease.

Conclusion

It is clear that the administration of food colouring as Ponceau 4R to rats caused many disturbance in different parameters of rats. It is recommended that the use of Ponceau 4R as food colouring must be limited due to its hazardous effect to the children and human health. It is recommended to use vitamin E as antioxidant to prevent the toxic effect of food colouring substance.

References

1. Abd El-Dayem SM. (1992): Toxic effect of tartrazine and amaranth on liver function of rats. *Proc. Egypt. Acad. Sci.*, 42: 133-138.
2. Abd El-Dayem SM. (2002): Biochemical alteration of pregnant albino rats intoxicated with dimethyl nitrosamine (DMNA). *J. Egypt. Ger. Soc. Zool.*, 3 (A) *Comparative Physiology*: 243-264.
3. Abd El-Rahiem AA, Yassin M, *et al.* (1999): Effect of sodium nitrite administration on acetylcholinesterase activity in discrete brain regions of rabbit. *J. Egypt. Ger. Soc. Zool.*, 30 (A) *Comparative Physiology*: 21-33.
4. Abu El-Zahab HSH, El-Khyat ZA, Sidhom G, *et al.* (1997): Physiological effects of some synthetic food colouring additives on rats. *Boll. Chem. Farm.*, 136 (10): 615-627.
5. Ahmed HH and Mona F. (2000): Protective effect of vitamin C and E against the toxic action of drinking sodium nitrate contaminated water in adult male albino rats. *J. Egypt. Ger. Soc. Zool.*, 32 (A) *Comparative Physiology*: 165-185.
6. AL-Sharkawi IM, Mansour MA, and El-Fiky BK. (1996): Non-permitted food

- colorants in the food products sold in the local markets in tanta city and the physiological derangements of their short term oral intake on liver and kidney functions. J. Union Arab Biol. Cairo, Zool., 6 (a): 351-371.
7. Amr TA, Budawy ME, *et al.* (1994): Effect of curacron toxicity on some liver functions.3-lipid metabolism and metabolic products.J.Union Arab Biol.,2(A): 263-282.
 8. Armitage P (1974): Statistical methods in medical research. Blackwell scientific Pub Oxford, London., PP: 116-120.
 9. Bagchi K and Puri S (1998): Free radicals and antioxidants in health and disease. Eastern Mediterranean Health J.,4 (2):350-356.
 10. Bär F & Griepentrog F (1960): Die Allergenwirkung von Fremden Stoffen in der Lebensmitteln, Med. u. Ernähr., 1, 99-104.
 11. Brantom PG, Stevenson BI, And Ingram AJ. (1987 a): A three generation reproduction study of Ponceau 4R in the rats. Food Chemical Toxicology, 25 (12): 963-968.
 12. Brantom PG, Stevenson BI, And Ingram AJ. (1987 b): Long term toxicity study of Ponceau 4R in rats using animals exposed in utero. Food and Chem. Toxicol., 25 (12): 955-962.
 13. Dinman BD, Hamdi EA, *et al.* (1963): Carbon tetrachloride (CCl₄) toxicity. III. Hepato structural and enzymatic change. Arch. Environ. Health, 7:630-646.
 14. Doumas BT. (1975): Colorimetric determination of total protein in serum or plasma. Clin. Chem., 21 (8): 1159-1166.
 15. Doumas BT, Watson WA, and Biggs HG. (1971): Albumin standards and measurements of serum albumin bromocresol green. Clin. Chem. Acta., 31: 87-96.
 16. Eder K, and Kirchgessner M. (1998): The effect of dietary vitamin E supply and moderately oxidized oil on activities of hepatic lipogenic enzymes in rats. Lipids, 33 (3): 277-283.
 17. Farag R, Abu-Raia S, *et al.* (1991): Safety evaluation of thyme and clove essential oils as natural antioxidant. AJAS., 18 (1):570-578.
 18. FAO & WHO (1991): Codex general standard for labeling of food additives when sold as such (World Wide Standard). Codex Alimentarius, 91: 51-91.
 19. Gaunt, I. F. *et al.* (1967): Acute (mouse and rat) and short-term (rat) toxicity studies on Ponceau 4R, Food Cosmet. Toxicol.; 5: 187-194.
 20. Gautam D, Shinha RK, And Milne DR. (1986): Interaction of Ponceau 4R with copper and effect of feeding Ponceau 4R on iron metabolism. J. Food Sci. and Technol. India; 23 (66): 303-307.
 21. Gomaa GMA. (1995): Protective effect of phospholipids and some vitamins against insecticides in toxication in male rats. Ph D Thesis, Dep Zool. Fac. Sci., Ain Shams University.
 22. Gorun V, Proinov I, Baltescu V, *et al.* (1978): Modified Ellman procedure for assay of cholinesterase in crude enzymatic preparation. Anal. Biochem., 86: 324-326.
 23. Grant D, and Butler WH. (1989): Chronic toxicity of sodium nitrite in male F344 rat. Food and Chem. Toxicol., 27 (9): 565-571.
 24. Hassan AA, El-Kalili MM, *et al.* (1995): Changes in serum lipid profile and esterases of rats after sublethal daily doses of dimethoate. J. Egypt. Pub. Health Ass., 19 (3): 431-437.
 25. Helal EG, Zaahkounk SA, and Hassan AB (1997): Biochemical and hematological effects of 8-quinaldine dimethyl carbamate methiodide on albino rats. J. Egypt. Ger. Soc. Zool., 24(A) Comparative Physiology: 119-133.
 26. Howard W and Janice E. (1989): An investigation of acetylcholinesterase activity following a high level acute exposure of pordoxo. Pesticide Biochem. And Physiol., 33: 125-131.
 27. Hurkat BC (1977): Some biochemical studies in rabbits (*Oryctolagus cuniculus*) during a hundred day oral administration of dieldrin (HEOD). Indian J. Anim. Sci., 47 (10): 671-676.
 28. Ignatov M (1976): Changes in some blood biochemical indices in chickens poisoned with thimet and lebaycide. Veter. Inarnameditsinkii Nauki, 13: 14-20.
 29. Isliker H, Weiser H and Moser U. (1997): Stabilization of rat heart mitochondria by α -tocopherol in rats. Internat. J. Vit. Nutr. Res., 67: 91-94.
 30. Katzung BG. (1995): Basic and clinical pharmacology. A lange Medical Book, 6th Ed. Eisei shikensho. Hokoku., 112: 48-56.
 31. Knight JA, Anderson S, and Rawie JM. (1972): Chemical basics of sulfophosphovanillin reaction for estimating total serum lipid. Clin. Chem., 18: 199-202.
 32. Lentner C, Lentner C, and Wink A (1982): International medical and pharmaceutical information. Vol 2, Ciba-Geigy limited, Basal, Switzerland.
 33. Luckens MM, and Phelps KI. (1969): Serum enzyme patterns in acute poisoning with organochlorine insecticides. J. Pharm. Sci., 58: 569-575.

34. Martinek RG (1970): Measurements of serum total cholesterol in practical clinical biochemistry. Vol. 1. 5th Ed. Pp: 655-657.
35. Mason PL, Gaunt IF, Butterwoeth KR, *et al.* (1974): Long term toxicity studies of carmosine in mice. Food and Cosm. Toxicol., 12 (516): 601-607.
36. Mcdoniel CD, Freed M, Balog JM, *et al.* (1993): Response of layer breeders to dietary acetylsalicylic acid. 4. Egg residue studies. Poultry Sci., 72: 1109-1117.
37. Mekawy IA, Hussein SY, *et al.* (1988): comparative studies on the effect of herbicide atrazine on some blood constituents and protein electrophoretic proteins of *Oreochromis niloticus* and *Chrysichthys quratus* at Assiut. J. Egypt. Ger. Soc. Zool., 19 (A) Comparative Physiology: 283-319.
38. Metin g, Atukeren P, Gumustas M, Belce A, and Kayserilioglu A. (2002): The effect of vitamin E treatment on oxidative stress generated in Trained rats. Tohoku J. Exp. Med., 198 (2): 47-53.
39. Meyer O, Kristiansen E, and Wurtzen G. (1989): Effect of dietary protein and butylated hydroxytoluene on the kidney of rats. Institute of Toxicol. Arch. Toxicol., 355 (suppl I): 5443.
40. Mitruka BM, Rawnsley HM, and Vadehra BV. (1977): Clinical biochemical and hematological reference values in normal experimental animals. Masson publishing USA, Inc. New York, Paris, Barcelona, Milan. Pp: 41-57.
41. Moriguchi S, Kobayashi N, and Kishino Y. (1990): High dietary intakes of vitamin E and cellular immune functions in rats. J. Nutr., 120 (9): 1069-1102.
42. Naziroglu M (1999): Protective role of intraperitoneally administered vitamin E and selenium in rats anesthetized with enflurane. Biol. Trace Elem. Res., 69 (3): 199-209.
43. Reitman S and Frankel S (1957): A colourimetric method for determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. Am. J. Clin. Path., 28: 56-61.
44. Rodak LC (1995): Determination of Hct value in routing testing in hematology in diagnostic hematology. W.B. saunders company, Philadelphia, London, Toronto. PP 128-144
45. Said HN, Sadhu HG, Bhatnagar VK, *et al.* (1992): Cardiac toxicity following short term exposure to methomyl in spray men rabbits. Human and Exp. Toxicol., 11 (2): 93-97.
46. Saleh SH. (1986): Biochemical studies on laboratory rats exposed to some food colours used in Egypt. M. Sc. Thesis. Fac. Agric., Cairo University.
47. Saleh F (1990): Effect of synthetic pyrethroid cypermethrin on lipid metabolism in pigeons. J. Egypt. Ger. Soc. Zool., 21 (A) Comparative Physiology: 67-77.
48. Shakoori *et al.* (1988): Effect of six months feeding on cypermethrin on the blood and liver of albino rats. J. Biochemical Toxicol., 3: 59-72.
49. Soheir LS, Khadiga G, and Soheir OE. (1996): Evaluation of serum ferritin in Egyptian children with protein energy malnutrition. J. Egypt. Ger. Soc. Zool., 21(A) Comparative Physiology: 1-16.
50. Soliman FA, Soliman MK, Abdo MS, *et al.* (1973): Respiration and circulation. In: veterinary Physiology, volume 1, 1st Ed. El-Kilany press.
51. Stevenson B I *et al.* (1980): Multi-generation toxicity study in rats with Ponceau 4R, Vol. I. Unpublished report No. 1/205/80, from the British Industrial Biological Research Association, submitted to WHO by the UK Colours Steering Group.
52. Stevenson B I *et al.* (1981): Long-term study in rats with Ponceau 4R using animals exposed in utero, Vol. I. Unpublished report No. 293/2/81, from the British Industrial Biological Research Association, submitted to WHO by the UK Colours Steering Group.
53. Tan YS, Nambiar R, and Yo SL. (1992): Prevalence of protein calorie malnutrition in general surgical patients. Ann. Acad. Med. Singapore, 21 (3): 334-338.
54. Trinder P (1969): Determination of glucose in blood using glucose oxidase with an alternative acceptor. Ann. Clin. Biochem., 6: 24-27.
55. Uchida K, Yasuhary N, Haruto T, *et al.* (1990): Effect of vitamin C depletion on serum cholesterol and lipoprotein levels in rats unable to synthesize ascorbic acid. J. Nutr., 120 (10): 1140-1148.
56. Van Kampen EJ and Zijlstra WG (1961): Standardization of Hæmoglobin-cyanide method. Clin Chem Acta 6: 538.
57. Wills JH. (1972): The measurement and significance of changes in the cholinesterase activities of erythrocytes and plasma in man and animals. Crit. Rev. Toxicol., 1: 153-160.
58. Yani M, Abd El-Dayem SM, and Abd El-Azim BH. (1991): Biochemical and histological changes due to preservative in rats. Egypt. J. Histol., 14 (2): 431-449.

دراسات كيميائية حيوية على تأثير معاملة بونسيية 4R و/ أو فيتامين اي (E) على ذكور الفئران

إيمان هلال*، محمود القصاص*، غادة زع سليمان**، سمير زيتون*، هشام عبد الواحد**
* قسم علم الحيوان-كلية العلوم-جامعة الأزهر (البنات)-** المعهد القومي للتغذية

المقدمة:

نظرا للتطور الهائل في هذه الايام فان استخدام المواد الملونة بونسيية 4R في الصناعات الغذائية جعل شكل الغذاء اكثر جاذبية للاطفال ، لذا تهدف هذه الدراسة لبيان تأثير بونسيية 4R (مواد ملونة) وبيان التأثير المتبادل لفيتامين اي (E) (مضاد للأكسدة طبيعي) على الفئران.

المواد والطرق:

تم قياس بعض المتغيرات مثل نسبة التغير في وزن الجسم، درجة حرارة الجسم، معدل ضربات القلب/دقيقة، عدد كرات الدم الحمراء والبيضاء، الهيموجلوبين، الهيماتوكريت، الدهون الكلية، الكوليسترول، البروتين الكلى، الأليومين، الجلوكوز، ألانين ترانس امينيز، اسبرتات ترانس امينيز، الكولين استيريز.

النتيجة:

وجد نقص معنوي في معظم المتغيرات مثل وزن الجسم، معدل التنفس حتى الشهر الثالث ثم اخذ في الزيادة المعنوية حتى الشهر السادس. وجد نقص معنوي في عدد كرات الدم البيضاء والكوليسترول والليبيدات الكلية في الدم والكبد، الجلوكوز، الكولين استيريز. بينما لوحظ ارتفاع معنوي في كوليسترول الدم، الليبيدات الكلية في العضلات، البروتين الكلى، الأليومين، ألانين ترانس امينيز، اسبرتات ترانس امينيز.

الاستنتاج:

نتيجة للتأثيرات العشوائية لمضافات الأغذية مثل بونسيية 4R ، فإنه يوصى بتحديد استخدام بونسيية 4R مع استخدام فيتامين اي (E) كمادة مضادة للأكسدة لمنع التأثير الضار.